FOR ANIMAL TREATMENT ONLY



PLEASE READ THE LABEL COMPLETELY BEFORE USE

Hazard Classification:

6.1D, 6.5B, 6.9B, 9.3C

Safety Directions

May be harmful if swallowed.
Wash hands and exposed skin before
meals and after use. Repeated exposure
may cause skin allergy. Avoid skin contact.
Triclabendazole may possibly affect the liver.
Handle with care.

First Aid

If swallowed DO NOT induce vomiting. For advice contact the National Poisons Centre 0800 POISON (0800 764766) or a doctor immediately. If skin or hair contact occurs remove contaminated clothing and flush skin and hair with running water.

Environmental Protection

Harmful to terrestrial vertebrates. Avoid release to the environment.

Emergency Response

In case of spillage wear appropriate protective clothing and prevent material from entering waterways. Absorb spills with inert material and place in waste containers. Wash the area with water and absorb with further inert material. Dispose of waste safely.

Disposal

Preferably dispose of the product by use. Otherwise dispose of product and packaging at an approved landfill or other approved facility. Triple rinse container with water. Crush or puncture and bury in a suitable landfill. DO NOT use container for any other purpose.

*For further details see the Safety Data Sheet.

FASINEX 24 can be given safely to all cattle including pregnant, young and stressed stock.

DRENCHING PROGRAMME

Immature liver fluke cause extensive damage when migrating through the liver tissue. Because FASINEX 24 is highly effective against immature as well as mature liver fluke, it should be given early in the season to reduce liver damage and prevent build up in the fluke population. Treatment should be repeated at 8 week intervals as necessary. Dairy cows should be treated only during the dry period, prior to calving.

Correct drenching technique should be used.

NET CONTENTS: 5 L



Elanco, Fasinex and the diagonal bar are trademarks owned or licensed by Eli Lilly and Company, its subsidiaries or affiliates.

Cattle



Cattle Dose weight (mL)

Up to 100 kg 4.5 mL 101–150 kg 6.5 mL

151-200 kg 8.5 mL

201-250 kg 10.5 mL 251-300 kg 15 mL

301-350 kg 17 mL

351-400 kg 19 mL 401-450 kg 22.5 mL

451–500 kg 25 mL

501-550 kg 27.5 mL 551-600 kg 30 mL

601-650 kg 32.5 mL

Cattle heavier than 650 kg should be dosed at 2.5 mL per 50 kg.

HELPLINE:

Phone 0800 ELANCO (0800 352 626)



It is an offence for users of this product to cause residues exceeding the relevant MRL in the New Zealand (Maximum Residues Limits of Agricultural Compounds) Food Standards.

WITHHOLDING PERIODS

1. MEAT:

Animals producing meat and offal for human consumption must not be slaughtered within 28 days of treatment.

2. MILI

Milk intended for sale for human consumption must be discarded for 35 days following treatment. NOT to be used in cows within 28 days of calving.

EFFICACY SPECTRUM

FASINEX 24 is a ready to use aqueous suspension of triclabendazole which is highly effective against early immature, immature and mature stages of liver fluke (Fasciola heaatica) in cattle.

DIRECTIONS

FASINEX 24 given as an oral drench and is suited for most types of automatic drenching gun.

1. SHAKE WELL BEFORE USE.

- 2. Give full dose orally according to dosing schedule.
- 3. A representative sample of animals should be weighed before treatment.
- Dose the herd to the heaviest animal by liveweight in each group. DO NOT underdose.
- Where there is a large variation in size within the group, dose rate should be based on the label directions for each weight range. Drafting into two or more lines may be appropriate, to avoid excessive overdosina.

By law the user must take due care, obtaining expert advice when necessary, to avoid unnecessary pain and distress when using this product other than as directed on the label.

Store in a cool, dry place out of direct sunlight, avoiding temperatures below 0° and above $35^{\circ}\text{C}.$

CONDITIONS OF SALE

The manufacturer of this animal remedy extends/grants to the purchaser a warranty that this animal remedy is reasonably fit for the purposes for which its use is recommended, provided that the purchaser uses the remedy only for the purpose for which it is recommended him and strictly in accordance with the directions on this container.

Registered pursuant to the ACVM Act 1997, No. A008127.

See <u>www.foodsafety.govt.nz</u> for registration conditions. Registered to Elanco Animal Health

(Division of Eli Lilly and Company (NZ) Limited), Level 1, 123 Ormiston Road, Botany Junction, Auckland 2016. Approved pursuant to the HSNO Act 1996, Approval Code: HSR001813.

See www.epa.govt.nz for approval conditions.

BATCH No: EXPIRY:

YL401385NZB 916779