

Literatuur



[Diagnostic accuracy and user-friendliness of 5 point-of-care D-dimer tests for the exclusion of deep vein thrombosis.](#)

Geersing GJ, Toll DB, Janssen KJ, Oudega R, Blikman MJ, Wijland R, de Vooght KM, Hoes AW, Moons KG. Clin Chem. 2010 Nov;56(11):1758-66. doi: 10.1373/clinchem.2010.147892.

[The cost-effectiveness of point-of-care D-dimer tests compared with a laboratory test to rule out deep venous thrombosis in primary care.](#)

Hendriksen JM, Geersing GJ, van Voorthuizen SC, Oudega R, Ten Cate-Hoek AJ, Joore MA, Moons KG, Koffijberg H. Expert Rev Mol Diagn. 2015 Jan;15(1):125-36. doi: 10.1586/14737159.2015.976202.

[Excluding venous thromboembolism using point of care D-dimer tests in outpatients: a diagnostic meta-analysis.](#)

Geersing GJ, Janssen KJ, Oudega R, Bax L, Hoes AW, Reitsma JB, Moons KG. BMJ. 2009 Aug 14;339:b2990. doi: 10.1136/bmj.b2990.

[Safely ruling out deep venous thrombosis in primary care.](#)

Büller HR, Ten Cate-Hoek AJ, Hoes AW, Joore MA, Moons KG, Oudega R, Prins MH, Stoffers HE, Toll DB, van der Velde EF, van Weert HC; AMUSE (Amsterdam Maastricht Utrecht Study on thromboEmbolism) Investigators..

Ann Intern Med. 2009 Feb 17;150(4):229-35

[Safe exclusion of pulmonary embolism using the Wells rule and qualitative D-dimer testing in primary care: prospective cohort study.](#)

Geersing GJ, Erkens PM, Lucassen WA, Buller HR, Cate HT, Hoes AW, Moons KG, Prins MH, Oudega R, van Weert HC, Stoffers HE.

BMJ. 2012 Oct 4;345:e6564. doi: 10.1136/bmj.e6564.

[NHG standaard 'Diep veneuze trombose en longembolie.](#)

Jip de Jong, Wim A.M. Lucassen, Geert-Jan Geersing, Jako S. Burgers en Tjerk Wiersma.

NHG-standaard 'Diepe veneuze trombose en longembolie'. Ned Tijdschr Geneesk. 2015;159:A8657